



« *They are satisfied,
They testify* »

FOCUS ON AGROBIO EUROPE

230 000 organic layers raised in open AIR divided in premises from 2 to 3 sections of 3 000 layers

50 000 « label rouge » layers raised in open air

Every breeding works under eggs takeover contract, There is no integration.

ALPHATECH offers in its range TONIFARM Poultry several alternatives adapted to breeding issues and usable in organic farming: **TONIFARM CI**, **TONIFARM P**, TONIFARM H (hepatoprotector), TONIFARM V (parasitism), TONIFARM AC (coccidiosis).

Denis PATUREL, ABE Manager:

« "Since 2009 we propose the **TONIFARM CI** for **IB** symptoms, and we successfully tried the **TONIFARM P** in 2013 with a breeder to reinforce the natural defenses of his layers. With 4 breeders in 2014 and with most of our breeders in 2015." »

Some testimonies:

Aurélie SAINTILLAN — 2 X 3 000 Organic layers



The first user of the TONIFARM CI with ABE in May 2009.

This breeding is located in an area, where broilers and layers concentration is very high. Very satisfied of the TONIFARM CI she decides to use the TONIFARM P in order to improve natural defenses of her animals from birth.

Thus, she uses the TONIFARM P from the chickens arrival 2 non-consecutive days per week and continues during the laying periode. Find below the results after an entire flock with this protocole and a 2nd flock still in progress in Week 43.

"After having received 2 consecutive flocks of heterogeneous pullets at 18 weeks during the past (flocks 1 and 2 : 1,400 and 1,300 Kg), I decided to set up (for the flocks 3 and 4) the TONIFARM P from the chickens arrival; This is when we have to start because I noted that the pullets arrive in breeding in good health and are more resistant in their new environment.

During BI passages, I still use the TONIFARM CI, laying falls are weaker than in the past. I don't use any booster vaccine during the laying period."



Technical results in August 2014 Use of TONIFARM P from chickens arrival.

Flock 3:

Pullets weight at 17 weeks: 1.400 Kg.
At 70 weeks: mortality 6% and 304 eggs/layer


Flock 4 : (in progress, arrived the 17/09/14)

Pullets weight at 18 weeks: 1.620 Kg
« Since the end of september, I used only twice the TONIFARM CI »
Flock results on the 20/03/15, week 43:
Mortality 1.5%, 148 eggs/layer and 93.2% of laying rate.

« I appointed that since I use the TONIFARM P, from brooding, layers are more resistant and that makes my job easier. »

Alain FLEURY — 2 sections of 3 000 Organic layers

« In the past, during IB passages, we could use 5 to 6 times the TONIFARM CI by flock. On the last flock, that just finished in march 2015, we used upon ABE advices the TONIFARM P 2 non-consecutive days per week (Monday and Wednesday) from the pullets arrival and during the entire round. »



IB at the week 51:
Laying decrease from 94% to 82% and back to 91.7% at the week 54.

Results of the flock at 70 weeks:
316 eggs per layer mortality 3.5%.

Only 2 IB passages for this flock.

« With the TONIFARM P distributed as preventive, during IB passages, instead of recovering in one month, layers recover in 15 days with the TONIFARM CI. »

Hervé PERROT — « 2 label rouge » henhouses 5 000 and 6 000 layers

« Since the flock beginning, on ABE technician advices, I distribute the TONIFARM P 2 non-consecutive days per week (Monday/Thursday). Compared to previous flocks, I have less IB passages. As soon as I see some IB symptoms, I double the dosage of TONIFARM P during 4 days running; For now, it works, knowing that if necessary, I have the TONIFARM CI. »

My results:

In week 60, my 6 000 henhouse is at 88% of laying rate and 2,20% of mortality.
My 5 000 henhouse : Despite an accident with a mortality of 500 layers, the laying rate is at 82% and layers are in good health.

« For me, the TONIFARM P acts like a fortifier to improve layers vitality »